

t13_wellord1 (TMd-
BrabC5gxu1FWHELQPzwGie9g4dULhWg2)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relat_1 : \iota \Rightarrow \iota$ be given. Let $k2_wellord1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X0 \in k1_relat_1 (k2_wellord1 X2 X1)) \Rightarrow ((X0 \in k1_relat_1 X2) \wedge (X0 \in X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k1_relat_1 (k2_wellord1 X1 X0)) (k1_relat_1 X1)) \wedge (r1_tarski (k1_relat_1 (k2_wellord1 X1 X0)) X0))$$