

t13_xreal_1

(TMYNs12QEfEhhJUqqM89h81tXppyi7QpSM9)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_xxreal_0 (k6_xcmplx_0 \\ & X0 X2) (k6_xcmplx_0 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k6_xcmplx_0 X0 X1) X2) \Rightarrow (r1_xxreal_0 \\ & (k6_xcmplx_0 X0 X2) X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (\\ & r1_xxreal_0 X0 X2)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 \\ & (k6_xcmplx_0 X0 X1)) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 X0 X3) (\\ & k6_xcmplx_0 X1 X2)))))) \end{aligned}$$