

t144_xreal_1 (TMKCZbTLfY-
fAu9K8CGhHz2CgTBFQScn5aie)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 k6_numbers X1)) \Rightarrow (r1_xxreal_0 (k7_xcmplx_0 X0 X1) k6_numbers))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 X1 k6_numbers)) \Rightarrow (r1_xxreal_0 (k7_xcmplx_0 X0 X1) k6_numbers))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 k6_numbers X1) \wedge (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X1)))))) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 (k7_xcmplx_0 X0 X1) k6_numbers) \wedge ((\neg(\neg r1_xxreal_0 X1 k6_numbers) \wedge (\neg r1_xxreal_0 X0 k6_numbers)) \wedge (\neg(\neg r1_xxreal_0 k6_numbers X1) \wedge (\neg r1_xxreal_0 k6_numbers X0))))))$$