

t146_xboolean
(TMYybHknHvjTDwXKhKeisZJRuzpLq3Lbw8j)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k9_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboolean : \iota$ be given. Let $k6_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboolean : \iota$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k6_xboolean X0 (k8_xboolean X0 X1) = k8_xboolean X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k6_xboolean X0 X1) = k1_xboolean)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k9_xboolean X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k8_xboolean X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Leftrightarrow ((X0 = k1_xboolean) \vee (X0 = k2_xboolean)) \quad (5)$$

Assume the following.

$$k2_xboolean = np_1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (k8_xboolean X0 X1 = k8_xboolean X1 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k8_xboolean X0 X1) = k1_xboolean))$$