

t146\_xreal\_1 (TMRDFuyDYge-  
Was6xefQHqW54E1eWFE1bNnf)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 (k2\_xcmplx\_0 X1 np\_1) X0))) \quad (1)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (r1\_xxreal\_0 X0 X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k6\_xcmplx\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X2) X1)))) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (6)$$

**Theorem 1**  $\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\neg r1\_xxreal\_0 X0 (k6\_xcmplx\_0 X0 np\_1)).$