

t148_xreal_1
(TMWpGVFua7UaBaSDKoj45iL8xEdogC2iTE4)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X1 (k4_xcmplx_0 X0)) \wedge (r1_xxreal_0 (k2_xcmplx_0 X1 X0) k6_numbers))) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (2)$$

Assume the following.

$$k2_xcmplx_0 np_1 (k4_xcmplx_0 np_1) = np_0 \quad (3)$$

Assume the following.

$$k2_xcmplx_0 np_1 (k4_xcmplx_0 np_1) = k6_numbers \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg(\neg r1_xxreal_0 X0 (k4_xcmplx_0 np_1)) \wedge (r1_xxreal_0 (k2_xcmplx_0 np_1 X0) k6_numbers))$$