

t14_arytm_1
(TMP56SDH2erngxjbP4YANqHUb9ewL9fNw6b)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_arytm_2 : \iota$ be given. Let $r1_arytm_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_arytm_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_arytm_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_arytm_2) \Rightarrow (\forall X2.(m1_subset_1 X2 k2_arytm_2) \Rightarrow (((r1_arytm_2 \\ & X0 X1) \wedge (r1_arytm_2 X2 X0)) \Rightarrow (k1_arytm_1 X1 (k1_arytm_1 X0 X2) = k7_arytm_2 \\ & (k1_arytm_1 X1 X0) X2)))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_arytm_2) \Rightarrow (\forall X2.(m1_subset_1 X2 k2_arytm_2) \Rightarrow (((r1_arytm_2 \\ & X0 X1) \wedge (r1_arytm_2 X2 X0)) \Rightarrow (k7_arytm_2 (k1_arytm_1 X1 X0) X2 = k1_arytm_1 \\ & X1 (k1_arytm_1 X0 X2)))) \end{aligned}$$