

t14_bvfunc_3

(TMdtdowrHN09U1a5DpGYdNFuvtXxVirYs6J)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_2 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_partit1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\
 & \quad (k1_bvfunc_2 X0))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\
 & \quad X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & \quad X0 k6_margrel1)))))) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge ((v1_funct_2 \\
 & \quad X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & \quad X0 k6_margrel1)))))) \Rightarrow (\forall X4. (m1_eqrel_1 X4 X0) \Rightarrow (r1_bvfunc_1 \\
 & \quad X0 (k9_bvfunc_1 X0 X2 X3) (k9_bvfunc_1 X0 (k6_bvfunc_2 X0 X2 X1 X4) \\
 & \quad (k7_bvfunc_2 X0 X3 X1 X4))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\
 & \quad (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & \quad k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge \\
 & \quad ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & \quad (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge \\
 & \quad ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & \quad (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (((r1_bvfunc_1 X0 X1 X2) \wedge (\\
 & \quad r1_bvfunc_1 X0 X2 X1)) \Rightarrow (r2_funct_2 X0 k6_margrel1 X1 X2)) \wedge (((r1_bvfunc_1 \\
 & \quad X0 X1 X2) \wedge (r1_bvfunc_1 X0 X2 X3)) \Rightarrow (r1_bvfunc_1 X0 X1 X3))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k1_partit1 X0))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 k6_margrel1)))) \Rightarrow (\forall X3.(m1_eqrel_1 X3 X0) \Rightarrow (r1_bvfunc_1 \\ X0 (k6_bvfunc_2 X0 X2 X1 X3) X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.k1_bvfunc_2 X0 = k1_partit1 X0 \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 k6_margrel1)))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k9_bvfunc_1 X0 X1 X2) \wedge \\ (v1_funct_2 (k9_bvfunc_1 X0 X1 X2) X0 k6_margrel1) \wedge (m1_subset_1 \\ (k9_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 \\ X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))) \wedge ((m1_subset_1 \\ X2 (k1_zfmisc_1 (k1_bvfunc_2 X0)) \wedge (m1_eqrel_1 X3 X0)))) \Rightarrow ((v1_funct_1 \\ (k7_bvfunc_2 X0 X1 X2 X3) \wedge ((v1_funct_2 (k7_bvfunc_2 X0 X1 X2 X3) \\ X0 k6_margrel1) \wedge (m1_subset_1 (k7_bvfunc_2 X0 X1 X2 X3) (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 \\ X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))) \wedge ((m1_subset_1 \\ X2 (k1_zfmisc_1 (k1_bvfunc_2 X0)) \wedge (m1_eqrel_1 X3 X0)))) \Rightarrow ((v1_funct_1 \\ (k6_bvfunc_2 X0 X1 X2 X3) \wedge ((v1_funct_2 (k6_bvfunc_2 X0 X1 X2 X3) \\ X0 k6_margrel1) \wedge (m1_subset_1 (k6_bvfunc_2 X0 X1 X2 X3) (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ & \quad (k1_bvfunc_2 X0))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & \quad X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge ((v1_funct_2 \\ & \quad X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad X0 k6_margrel1)))))) \Rightarrow (\forall X4.(m1_eqrel_1 X4 X0) \Rightarrow (r1_bvfunc_1 \\ & \quad X0 (k6_bvfunc_2 X0 (k9_bvfunc_1 X0 X2 X3) X1 X4) (k9_bvfunc_1 X0 (\\ & \quad k6_bvfunc_2 X0 X2 X1 X4) (k7_bvfunc_2 X0 X3 X1 X4)))))) \end{aligned}$$