

t14_calcul_1 (TMYVN-
Mms8id5Z7TV7QwUWyGQvhNg7qRSBME)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 X2))) \Rightarrow ((X2 = k10_finseq_1 X0 X1) \Leftrightarrow ((k3_finseq_1 X2 = np_2) \wedge ((k1_funct_1 X2 np_1 = X0) \wedge (k1_funct_1 X2 np_2 = X1)))) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (4)$$

Assume the following.

$$r1_xxreal_0 np_2 np_2 \quad (5)$$

Assume the following.

$$r1_xxreal_0 \ np_1 \ np_2 \tag{6}$$

Assume the following.

$$r1_xxreal_0 \ np_1 \ np_1 \tag{7}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 \ X1 \ X0) \Leftrightarrow (m1_finseq_1 \ X1 \ X0) \tag{8}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{9}$$

Assume the following.

$$\forall X0.((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow (k4_finseq_1 \ X0 = k9_xtuple_0 \ X0) \tag{10}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 \ X1) \wedge (v4_relat_1 \ X1 \ X0)) \Rightarrow (k1_relset_1 \ X0 \ X1 = k9_xtuple_0 \ X1) \tag{11}$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 \ (k10_finseq_1 \ X0 \ X1)) \wedge (v1_funct_1 \ (k10_finseq_1 \ X0 \ X1)) \tag{12}$$

Assume the following.

$$\forall X0.\forall X1.v1_finseq_1 \ (k10_finseq_1 \ X0 \ X1) \tag{13}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 \ X1 \ X0) \Rightarrow ((v1_funct_1 \ X1) \wedge (v1_finseq_1 \ X1) \wedge (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k5_numbers \ X0)))) \tag{14}$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 \ X1 \ X0) \Rightarrow ((v1_relat_1 \ X1) \wedge (v1_funct_1 \ X1) \wedge (v1_finseq_1 \ X1)) \tag{15}$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \wedge ((v1_relat_1 \ X1) \wedge ((v1_funct_1 \ X1) \wedge (v1_finseq_1 \ X1)))) \Rightarrow ((v1_relat_1 \ (k7_finseq_1 \ X0 \ X1)) \wedge ((v1_funct_1 \ (k7_finseq_1 \ X0 \ X1)) \wedge (v1_finseq_1 \ (k7_finseq_1 \ X0 \ X1)))) \tag{16}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\
& \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\
& \quad X1))) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 \\
& \quad X2)))) \Rightarrow ((X2 = k7_finseq_1 X0 X1) \Leftrightarrow ((k4_finseq_1 X2 = k2_finseq_1 \\
& \quad (k2_nat_1 (k3_finseq_1 X0) (k3_finseq_1 X1))) \wedge ((\forall X3.(\\
& \quad v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X0) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 \\
& \quad X0 X3))) \wedge (\forall X3.(v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X1) \Rightarrow \\
& \quad (k1_funct_1 X2 (k2_nat_1 (k3_finseq_1 X0) X3) = k1_funct_1 X1 X3))))))))) \quad (17)
\end{aligned}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (18)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\
& ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge \\
& \quad (v1_finseq_1 X0)))) \quad (19)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1. \forall X2. \forall X3. \\
& (m2_finseq_1 X3 (k3_qc_lang X0)) \Rightarrow ((np_1 \in k1_relset_1 k5_numbers \\
& (k10_finseq_1 X1 X2)) \wedge ((np_2 \in k1_relset_1 k5_numbers (k10_finseq_1 \\
& X1 X2)) \wedge ((k1_funct_1 (k7_finseq_1 X3 (k10_finseq_1 X1 X2)) (k2_nat_1 \\
& (k3_finseq_1 X3) np_1) = X1) \wedge (k1_funct_1 (k7_finseq_1 X3 (k10_finseq_1 \\
& X1 X2)) (k2_nat_1 (k3_finseq_1 X3) np_2) = X2))))))
\end{aligned}$$