

t14_cqc_the3
(TMU9utBoES73oV5R6LoSiovtzyYTFxibUMZ)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $r2_cqc_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_subset_1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k4_cqc_the1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow (r2_cqc_the3 X0 X1 (k4_cqc_the1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.m1_subset_1 (k1_subset_1 X0) (k1_zfmisc_1 X0) \quad (2)$$

Theorem 1

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (r2_cqc_the3 X0 (k1_subset_1 (k3_cqc_lang X0)) (k4_cqc_the1 X0))$$