

t14_fin_topo
(TMXnYyrUn5feRx8pmCYB4oQwm3g618C5739)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1)) \wedge (r1_xboole_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k9_fin_topo X0 X1 = ReplSep (toset (\lambda X2 : \iota. m1_subset_1 X2 (u1_struct_0 X0))) (\lambda X2 : \iota. \neg r1_xboole_0 (k1_fin_topo X0 X2) X1) (\lambda X2 : \iota. X2)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Theorem 1

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k9_fin_topo X0 X1) (k9_fin_topo X0 X2))))))$$