

## t14\_funct\_9

(TMM8MEpL724cXWT8KPxsBaLHSUHFQwpGeqS)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_9 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 \\ & X1) \wedge (v3\_valued\_0 X1))) \Rightarrow ((v1\_funct\_9 X1 X0) \Leftrightarrow ((X0 \neq k6\_numbers) \wedge \\ & (\forall X2.(v1\_xreal\_0 X2) \Rightarrow ((X2 \in k9\_xtuple\_0 X1) \Rightarrow ((k2\_xcmplx\_0 \\ & X2 X0 \in k9\_xtuple\_0 X1) \wedge ((k6\_xcmplx\_0 X2 X0 \in k9\_xtuple\_0 X1) \wedge (k1\_funct\_1 \\ & X1 X2 = k1\_funct\_1 X1 (k2\_xcmplx\_0 X2 X0)))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow (k2\_xcmplx\_0 X0 X1 = k6\_xcmplx\_0 X0 (k4\_xcmplx\_0 X1))) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 \\ & X1) \wedge (v3\_valued\_0 X1))) \Rightarrow ((v1\_funct\_9 X1 X0) \Rightarrow (\forall X2.(v1\_xreal\_0 \\ & X2) \Rightarrow ((X2 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 X1 X2 = k1\_funct\_1 X1 (k6\_xcmplx\_0 \\ & X2 X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow ((k4\_xcmplx\_0 X0 = k6\_numbers) \Rightarrow (X0 = k6\_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k2\_xcmplx\_0 X0 (k4\_xcmplx\_0 X1) = k6\_xcmplx\_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow ((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0)) \wedge (v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 \\ X1)) \Rightarrow ((v1\_funct\_9 X1 X0) \Leftrightarrow ((X0 \neq k6\_numbers) \wedge (\forall X2.(v1\_xreal\_0 \\ X2) \Rightarrow (((X2 \in k9\_xtuple\_0 X1) \Rightarrow (k2\_xcmplx\_0 X2 X0 \in k9\_xtuple\_0 X1)) \wedge \\ (((k2\_xcmplx\_0 X2 X0 \in k9\_xtuple\_0 X1) \Rightarrow (X2 \in k9\_xtuple\_0 X1)) \wedge \\ (X2 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 X1 X2 = k1\_funct\_1 X1 (k2\_xcmplx\_0 \\ X2 X0)))))))))) \quad (7) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k2\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (9)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v3\_valued\_0 X1))) \Rightarrow ((v1\_funct\_9 X1 X0) \Rightarrow (v1\_funct\_9 X1 (k4\_xcmplx\_0 X0))))$$