

t14_mcart_1 (TMFB-
NuyUw7TuXJX5rtei6jZmEEYQR5BTm1s)

October 27, 2020

Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X0 \in k2_zfmisc_1 X1 X2) \Rightarrow ((k1_xtuple_0 X0 \in X1) \wedge (k2_xtuple_0 X0 \in X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (X0 \in k2_zfmisc_1 (k1_tarski X1) (k1_tarski X2)) \Rightarrow ((k1_xtuple_0 X0 = X1) \wedge (k2_xtuple_0 X0 = X2))$$