

t14_measure6
(TMdojoyckbJi7zcfVDgeYUxSJgqBHAvKDuc)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v6_xxreal_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k7_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_supinf_2 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_xxreal_2 : \iota \Rightarrow \iota$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k1_xxreal_2 (k1_xxreal_1 X0 X1) = X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (k8_supinf_2 X0 = k1_xxreal_2 X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v2_membered (k1_xxreal_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k7_numbers) \Rightarrow (v1_xxreal_0 X0) \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge ((v6_xxreal_2 X0) \wedge (m1_subset_1 \\ & X0 (k1_zfmisc_1 k1_numbers)))) \Rightarrow (\forall X1.(m1_subset_1 X1 k7_numbers) \Rightarrow \\ & ((\exists X2.(m1_subset_1 X2 k7_numbers) \wedge ((r1_xxreal_0 X2 X1) \wedge \\ & (X0 = k1_xxreal_1 X2 X1))) \Rightarrow (X1 = k8_supinf_2 X0))) \end{aligned}$$