

t14_normsp_1 (TMHRC- CigLa3jBHpzWgR27AnPHkJqbVbCAtz)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow ((k9_xtuple_0 X0 = k1_xboole_0) \Leftrightarrow (k10_xtuple_0 X0 = k1_xboole_0)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 (k1_tarski X1)) \Leftrightarrow ((X0 = k1_xboole_0) \vee (X0 = k1_tarski X1)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\exists X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 k5_numbers (u1_struct_0 X0)) \wedge (m1_subset_1 X2 \\ & (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (u1_struct_0 X0)))))) \wedge \\ & (k10_xtuple_0 X2 = k6_domain_1 (u1_struct_0 X0) X1))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ \neg v2_struct_0 X1) \wedge (l1_struct_0 X1)) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (u1_struct_0 X1)) \Rightarrow (((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers \\ (u1_struct_0 X1)) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (u1_struct_0 X1)))))) \Leftrightarrow ((k9_xtuple_0 X0 = k5_numbers) \wedge \\ (\forall X3.(X3 \in k5_numbers) \Rightarrow (m1_subset_1 (k1_funct_1 X0 X3) \\ (u1_struct_0 X1))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (13)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers (u1_struct_0 X0)) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (u1_struct_0 \\ & X0)))))) \Rightarrow (\neg(\exists X2.(m1_subset_1 X2 (u1_struct_0 X0)) \wedge (\forall X3. \\ & (v7_ordinal1 X3) \Rightarrow (k1_funct_1 X1 X3 = X2))) \wedge (\forall X2.(m1_subset_1 \\ & X2 (u1_struct_0 X0) \Rightarrow (k10_xtuple_0 X1 \neq k6_domain_1 (u1_struct_0 \\ & X0) X2)))) \end{aligned}$$