

t14_topgen_1
(TMWdmu6xr2QZMSCnuDoyshNJH46MZVML8gq)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k2_tops_1 X0 X1 = k7_subset_1 (u1_struct_0 X0) (k2_pre_topc X0 X1) (k1_tops_1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v4_pre_topc X1 X0) \Leftrightarrow (k2_tops_1 X0 X1 = k7_subset_1 (u1_struct_0 X0) X1 (k1_tops_1 X0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(k4_xboole_0 X0 X1 = k1_xboole_0) \Leftrightarrow (r1_tarski X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1.(l1_pre_topc X1) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X1))) \Rightarrow (((v3_pre_topc X3 X1) \Rightarrow (k1_tops_1 X1 X3 = X3)) \wedge ((k1_tops_1 X0 X2 = X2) \Rightarrow (v3_pre_topc X2 X0)))))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (5)$$

Assume the following.

$$\forall X0.(l1_pre_topc\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0\ X0))) \Rightarrow (r1_tarski\ X1\ (k2_pre_topc\ X0\ X1))) \quad (6)$$

Assume the following.

$$\forall X0.(l1_pre_topc\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0\ X0))) \Rightarrow (r1_tarski\ (k1_tops_1\ X0\ X1)\ X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0)) \Rightarrow (k7_subset_1\ X0\ X1\ X2 = k4_xboole_0\ X1\ X2) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc\ X0) \wedge (m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow (m1_subset_1\ (k2_pre_topc\ X0\ X1)\ (k1_zfmisc_1\ (u1_struct_0\ X0))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1_tarski\ X0\ X1) \wedge (r1_tarski\ X1\ X0)) \quad (10)$$

Theorem 1

$$\forall X0.((v2_pre_topc\ X0) \wedge (l1_pre_topc\ X0)) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0\ X0))) \Rightarrow (((v3_pre_topc\ X1\ X0) \wedge (v4_pre_topc\ X1\ X0)) \Leftrightarrow (k2_tops_1\ X0\ X1 = k1_xboole_0)))$$