

# t14\_topreal6 (TMQxPZuE- BiZAG61KpwDYuRLw1ok91LS36tA)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k18\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k16\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k6\_xcmplx\_0 X1 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (k7\_real\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k18\_complex1 X0 = k16\_complex1 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (v1\_xreal\_0 (k6\_xcmplx\_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (m1\_subset\_1 (k18\_complex1 X0) k1\_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k16\_complex1 X0 = X0)) \wedge ((\neg r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k16\_complex1 X0 = k4\_xcmplx\_0 X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (7)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (((r1\_xxreal\_0 \\ & X0 X1) \wedge (r1\_xxreal\_0 X2 X3)) \Rightarrow (k7\_real\_1 (k18\_complex1 (k6\_xcmplx\_0 \\ & X1 X0)) (k18\_complex1 (k6\_xcmplx\_0 X3 X2)) = k2\_xcmplx\_0 (k6\_xcmplx\_0 \\ & X1 X0) (k6\_xcmplx\_0 X3 X2)))))) \end{aligned}$$