

t156_sheffer2

(TMZZ4pnEHismVWcFaj2bqBpD9hwdFZ85uCU)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v10_sheffer1 : \iota \Rightarrow o$ be given. Let $v11_sheffer1 : \iota \Rightarrow o$ be given. Let $v12_sheffer1 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ & X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (\forall X4. \\ & (m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X2 X4 = k5_sheffer1 \\ & X0 X2 (k5_sheffer1 X0 (k5_sheffer1 X0 X4 X1) (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 (k5_sheffer1 X0 X4 (k5_sheffer1 X0 X3 (k5_sheffer1 X0 X3 X3))) \\ & (k5_sheffer1 X0 X4 (k5_sheffer1 X0 X3 (k5_sheffer1 X0 X3 X3)))))))))) \\ & (1) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ & X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 \\ & X0 (k5_sheffer1 X0 X3 X2) (k5_sheffer1 X0 X3 (k5_sheffer1 X0 X1 (k5_sheffer1 \\ & X0 X1 X1))) = X3)))) \\ & (2) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((\neg v2_struct_0 X0) \wedge (l1_sheffer1 \\ & X0)) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 \\ & X0)))) \Rightarrow (m1_subset_1 (k5_sheffer1 X0 X1 X2) (u1_struct_0 X0)) \\ & (3) \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v10_sheffer1 X0) \wedge ((v11_sheffer1 \\ & X0) \wedge ((v12_sheffer1 X0) \wedge (l1_sheffer1 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 \\ & X0 X1 X3 = k5_sheffer1 X0 X1 (k5_sheffer1 X0 (k5_sheffer1 X0 X3 X2) \\ & (k5_sheffer1 X0 X3 X1)))))) \end{aligned}$$