

t15\_analoaf (TMLR-  
DoqW2C9d96wJFMDmraEfSZ4PmjYSpps)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_analoaf : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (3)$$

Assume the following.

$$\neg r1\_xxreal\_0 np\_1 np\_0 \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow \\
& (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 \\
& X3 (u1\_struct\_0 X0)) \Rightarrow (\forall X4.(m1\_subset\_1 X4 (u1\_struct\_0 \\
& X0)) \Rightarrow ((r1\_analoaf X0 X1 X2 X3 X4) \Leftrightarrow (\neg(X1 \neq X2) \wedge ((X3 \neq X4) \wedge (\forall X5. \\
& (m1\_subset\_1 X5 k1\_numbers) \Rightarrow (\forall X6.(m1\_subset\_1 X6 k1\_numbers) \Rightarrow \\
& (\neg(\neg r1\_xxreal\_0 X5 k6\_numbers) \wedge (\neg r1\_xxreal\_0 X6 k6\_numbers) \wedge \\
& (k1\_rlvect\_1 X0 (k5\_algstr\_0 X0 X2 X1) X5 = k1\_rlvect\_1 X0 (k5\_algstr\_0 \\
& X0 X4 X3) X6))))))))))))) \tag{6}
\end{aligned}$$

**Theorem 1**

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow \\
& (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 \\
& X3 (u1\_struct\_0 X0)) \Rightarrow (\forall X4.(m1\_subset\_1 X4 (u1\_struct\_0 \\
& X0)) \Rightarrow ((k5\_algstr\_0 X0 X1 X2 = k5\_algstr\_0 X0 X3 X4) \Rightarrow (r1\_analoaf \\
& X0 X2 X1 X4 X3)))))))))
\end{aligned}$$