

t15_fib_num3 (TMGWgwWaYVAdZh- Nev3kuPaaobqfoDY85sYu)

October 27, 2020

Let $k1_fib_num3 : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $np_4 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$k1_fib_num3\ np_2 = np_3 \tag{1}$$

Assume the following.

$$\begin{aligned} & (k1_fib_num3\ k6_numbers = np_2) \wedge ((k1_fib_num3\ np_1 = np_1) \wedge \\ & (\forall X0.(v7_ordinal1\ X0) \Rightarrow (k1_fib_num3\ (k2_nat_1\ (k1_nat_1 \\ & X0\ np_1)\ np_1) = k2_nat_1\ (k1_fib_num3\ X0)\ (k1_fib_num3\ (k1_nat_1 \\ & X0\ np_1)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_3) \wedge (m2_subset_1\ np_3\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_3\ k5_numbers) \wedge (m1_subset_1\ np_3\ k1_numbers)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_2) \wedge (m2_subset_1\ np_2\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_2\ k5_numbers) \wedge (m1_subset_1\ np_2\ k1_numbers)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_1) \wedge (m2_subset_1\ np_1\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_1\ k5_numbers) \wedge (m1_subset_1\ np_1\ k1_numbers)) \end{aligned} \tag{5}$$

Assume the following.

$$k2_xcmplx_0\ np_3\ np_1 = np_4 \tag{6}$$

Assume the following.

$$k2_xcmplx_0 \ np_2 \ np_1 = np_3 \quad (7)$$

Assume the following.

$$k2_xcmplx_0 \ np_1 \ np_1 = np_2 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 \ X0 \ k5_numbers)\wedge(v7_ordinal1 \ X1))\Rightarrow(k2_nat_1 \ X0 \ X1 = k2_xcmplx_0 \ X0 \ X1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 \ X0)\wedge(m1_subset_1 \ X1 \ k5_numbers))\Rightarrow(k1_nat_1 \ X0 \ X1 = k2_xcmplx_0 \ X0 \ X1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 \ X0 \ k5_numbers)\wedge(v7_ordinal1 \ X1))\Rightarrow(k2_nat_1 \ X0 \ X1 = k2_nat_1 \ X1 \ X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k4_ordinal1)\Rightarrow(v7_ordinal1 \ X0) \quad (13)$$

Theorem 1 $k1_fib_num3 \ np_3 = np_4$.