

t15_finseq_3

(TMPfxJZ3GkVLcfuQp1HQiz7i45TsXYjYJAN)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k2_xboole_0 (k2_finseq_1 X0) (k1_tarski (k1_nat_1 X0 np_1))) = k2_finseq_1 (k1_nat_1 X0 np_1) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_xboole_0 X0 X1) \Rightarrow (k4_xboole_0 (k2_xboole_0 X0 X1) X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xboole_0 (k2_finseq_1 X0) (k1_tarski (k1_nat_1 X0 np_1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(r1_xboole_0 X0 X1) \Rightarrow (r1_xboole_0 X1 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(m1_subset_1 (k2_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (12)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(k7_subset_1 k5_numbers (k2_finseq_1 (k1_nat_1 X0 np_1)) (k2_finseq_1 X0) = k1_tarski (k1_nat_1 X0 np_1))$$