

# t15\_finseq\_4 (TMPVGED- cazVe4ZDnXJi2mwLbHvQZTBKc3VN)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((X1 \in k1\_relset\_1 k5\_numbers X0) \Leftrightarrow \\ & ((r1\_xxreal\_0 np\_1 X1) \wedge (r1\_xxreal\_0 X1 (k3\_finseq\_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ((v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1.((v1\_relat\_1 X1) \wedge ((v5\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1))) \Rightarrow (\forall X2.(X2 \in k9\_xtuple\_0 X1) \Rightarrow (k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (7)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(\forall X1.\forall X2.(m2\_finseq\_1 X2 X1)\Rightarrow(((r1\_xxreal\_0 np\_1 X0)\wedge(r1\_xxreal\_0 X0 (k3\_finseq\_1 X2)))\Rightarrow(k7\_partfun1 X1 X2 X0 = k1\_funct\_1 X2 X0)))$$