

t15\_funct\_2 (TMK-  
wzVsYEhm4ezX8hGaPXgvMV4KCBcMm6P<sub>x</sub>)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. \\ & ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 \\ & (k3\_relat\_1 X1 X2) X0 = k1\_funct\_1 X2 (k1\_funct\_1 X1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow ( \\ & k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (((X1 \neq k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 \\ & X1) \Leftrightarrow (X0 = k1\_relset\_1 X0 X2))) \wedge ((X1 = k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 \\ & X2 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X3) \wedge \\ & ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))))) \Rightarrow (\forall X4. ((v1\_relat\_1 X4) \wedge (v1\_funct\_1 X4)) \Rightarrow (( \\ & X2 \in X0) \Rightarrow ((X1 = k1\_xboole\_0) \vee (k1\_funct\_1 (k3\_relat\_1 X3 X4) X2 = \\ & k1\_funct\_1 X4 (k1\_funct\_1 X3 X2)))))) \end{aligned}$$