

t15_funct_3
(TMWq1kKKq4jf3shQos7huCpLJs8MmMd9NmL)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_tarski : \iota \Rightarrow \iota$ be given. Let $k1_funct_3 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski \\ X0 (k1_zfmisc_1 (k9_xtuple_0 X1))) \Rightarrow (k7_relat_1 X1 (k3_tarski \\ X0) = k3_tarski (k7_relat_1 (k1_funct_3 X1) X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (v1_partfun1 X1 X0 \Leftrightarrow (k1_relset_1 X0 X1 = X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (\neg v1_xboole_0 X1) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v1_funct_2 X2 X0 X1) \Rightarrow (v1_partfun1 X2 X0))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (6)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.(\neg v1_xboole_0 X2)\Rightarrow(\forall X3. \\ ((v1_funct_1 X3)\wedge((v1_funct_2 X3 X0 X2)\wedge(m1_subset_1 X3 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X2))))))\Rightarrow((r1_tarski X1 (k1_zfmisc_1 X0))\Rightarrow(k7_relat_1 \\ X3 (k3_tarski X1) = k3_tarski (k7_relat_1 (k1_funct_3 X3) X1))))$$