

t15_funct_7

(TMSV2425GHmxVrcEVyed5m1PCJLHTqpVemy)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge \\ (v1_funct_1 X2)) \Rightarrow ((k9_xtuple_0 X1 = k9_xtuple_0 X2) \Rightarrow (k1_funct_4 \\ (k1_funct_4 X0 X1) X2 = k1_funct_4 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarski X1) \quad (6)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ \forall X3. k1_funct_4 (k1_funct_4 X0 (k7_funcop_1 X1 X2)) (k7_funcop_1 \\ X1 X3) = k1_funct_4 X0 (k7_funcop_1 X1 X3)) \end{aligned}$$