

t15_jordan3 (TMNuJE- jhrP6MhopMYgTgVuRAAdCBQ6ZbVmKg)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_jordan3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_5 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_topreal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & ((k1_funct_1 X0 np_1 = k1_funct_1 (k3_finseq_5 X0) (k3_finseq_1 \\ & X0)) \wedge (k1_funct_1 X0 (k3_finseq_1 X0) = k1_funct_1 (k3_finseq_5 \\ & X0) np_1)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow (k4_finseq_5 X0 X1 = k3_finseq_5 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow (k4_finseq_5 X0 (k4_finseq_5 X0 X1) = X1) \quad (4)$$

Assume the following.

$$\forall X0. ((v4_topreal1 X0) \wedge (m1_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (v4_topreal1 (k3_finseq_5 X0)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(m2_finseq_1 (k4_finseq_5 X0 X1) X0) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & ((r1_jordan3 X0 X1 X2)\Leftrightarrow((v4_topreal1 X0)\wedge((k1_funct_1 X0 np_1 = \\ & X1)\wedge(k1_funct_1 X0 (k3_finseq_1 X0) = X2)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & ((r1_jordan3 X0 X1 X2)\Rightarrow(r1_jordan3 (k4_finseq_5 (u1_struct_0 \\ & (k15_euclid np_2)) X0) X2 X1)))) \end{aligned}$$