

## t15\_matrix\_1

(TMQbH1bbALw6A8aodXedrQf2hjGZaVYyDZe)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_matrix\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k10\_xtuple\_0 X1 = k1\_tarski X0))) \quad (1)$$

Assume the following.

$$\forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (m2\_finseq\_1 X1 X0) \Rightarrow (m1\_matrix\_1 (k9\_finseq\_1 X1) X0 np\_1 (k3\_finseq\_1 X1))) \quad (2)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (4)$$

Assume the following.

$$\forall X0. v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (5)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 (k5\_finseq\_1 X0)) \wedge (v1\_funct\_1 (k5\_finseq\_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (7)$$

**Theorem 1**

$$\forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (m1\_matrix\_1 (k9\_finseq\_1 (k12\_finseq\_1 X0 X1)) X0 np\_1 np\_1))$$