

t15_mesfun6c (TMWj-
jAoCFeeibKPrfCNT7RNPmtMM174zA8r)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & \quad m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\\ & \quad \forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad X0 k2_numbers)))) \Rightarrow ((k1_relset_1 X0 (k2_valued_1 X0 k2_numbers \\ & \quad k2_numbers X1 X2) = k9_subset_1 X0 (k1_relset_1 X0 X1) (k1_relset_1 \\ & \quad X0 X2)) \wedge (\forall X3. (m1_subset_1 X3 X0) \Rightarrow ((X3 \in k1_relset_1 X0 (\\ & \quad k2_valued_1 X0 k2_numbers k2_numbers X1 X2) \Rightarrow (k7_partfun1 k2_numbers \\ & \quad (k2_valued_1 X0 k2_numbers k2_numbers X1 X2) X3 = k8_complex1 (k7_partfun1 \\ & \quad k2_numbers X1 X3) (k7_partfun1 k2_numbers X2 X3))))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X1) \wedge ((v1_prob_1 X1 X0) \wedge \\ & ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 \\ & X0)))))) \Rightarrow (\forall X2. (m1_prob_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((\neg v1_xboole_0 \\ & X1)\wedge((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\wedge((m1_subset_1 X2 X1)\wedge(m1_subset_1 X3 X1)))\Rightarrow \\ & (k5_prob_1 X0 X1 X2 X3 = k3_xboole_0 X2 X3) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((\neg v1_xboole_0 \\ & X1)\wedge((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\wedge((m1_subset_1 X2 X1)\wedge(m1_subset_1 X3 X1)))\Rightarrow \\ & (m1_prob_1 (k5_prob_1 X0 X1 X2 X3) X0 X1) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Leftrightarrow(\forall X1.\neg X1 \in X0) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0)\Rightarrow((m1_subset_1 X1 X0)\Leftrightarrow \\ & (X1 \in X0)))\wedge((v1_xboole_0 X0)\Rightarrow((m1_subset_1 X1 X0)\Leftrightarrow(v1_xboole_0 \\ & X1))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (12)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\forall X3.(\\ & (v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 \\ & k2_numbers)))) \Rightarrow (((k1_relset_1 X0 X2 \in X1) \wedge (k1_relset_1 X0 X3 \in \\ & X1)) \Rightarrow (k1_relset_1 X0 (k2_valued_1 X0 k2_numbers k2_numbers X2 \\ & X3) \in X1)))))) \end{aligned}$$