

t15_ordinal1 (TMH-
coS2QDZnwsWJH6QRgUDnvaui7m6fh9zS)

October 27, 2020

Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $r3_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_ordinal1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v3_ordinal1 X0) \wedge (v3_ordinal1 X1)) \Rightarrow (r1_ordinal1 X0 X1) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r3_xboole_0 X0 X1) \Leftrightarrow ((r1_tarski X0 X1) \vee (r1_tarski X1 X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v3_ordinal1 X0) \wedge (v3_ordinal1 X1)) \Rightarrow (r1_ordinal1 X0 X1) \vee (r1_ordinal1 X1 X0) \quad (3)$$

Theorem 1

$$\forall X0. (v3_ordinal1 X0) \Rightarrow (\forall X1. (v3_ordinal1 X1) \Rightarrow (r3_xboole_0 X0 X1))$$