

t15_tdlat_2

(TMHoHLbw69RLHHVPzY3UUdekYpgMZCkP5YJ)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pcomps_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\forall X2. (X2 \in X0) \Rightarrow (r1_tarski X2 X1)) \Rightarrow (r1_tarski (k3_tarski X0) X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (r1_tarski X0 (k3_tarski X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k2_pre_topc X0 X1) (k2_pre_topc X0 X2)))))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (k5_setfam_1 X0 X1 = k3_tarski X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k5_setfam_1 X0 X1) (k1_zfmisc_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc\ X0)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))))\Rightarrow(m1_subset_1\ (k1_pcomps_1\ X0\ X1)\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0))))\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0))))\Rightarrow((X2 = k1_pcomps_1\ X0\ X1)\Leftrightarrow(\forall X3.(m1_subset_1\ X3\ (k1_zfmisc_1\ (u1_struct_0\ X0))\Rightarrow((X3 \in X2)\Leftrightarrow(\exists X4.(m1_subset_1\ X4\ (k1_zfmisc_1\ (u1_struct_0\ X0))\wedge((X3 = k2_pre_topc\ X0\ X4)\wedge(X4 \in X1)))))))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} \forall X0.((v2_pre_topc\ X0)\wedge(l1_pre_topc\ X0)\Rightarrow(\forall X1. \\ (m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0))))\Rightarrow \\ (r1_tarski\ (k5_setfam_1\ (u1_struct_0\ X0)\ (k1_pcomps_1\ X0\ X1)) \\ (k2_pre_topc\ X0\ (k5_setfam_1\ (u1_struct_0\ X0)\ X1)))) \end{aligned}$$