

t15_weierstr
(TMXEEgjXdgVVGxuZQLuJjVhu5hs5jKs7Uds)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k3_topmetr : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_weierstr : \iota \Rightarrow \iota$ be given. Let $k7_reset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_seq_4 : \iota \Rightarrow \iota$ be given. Let $k1_weierstr : \iota \Rightarrow \iota$ be given. Let $k5_seq_4 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\
& X0))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (u1_struct_0 \\
& X0) (u1_struct_0 k3_topmetr)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
& k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 k3_topmetr)))))) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow \\
& (\neg (X2 \neq k1_xboole_0) \wedge ((v2_compts_1 X2 X0) \wedge ((v5_pre_topc X1 X0 \\
& k3_topmetr) \wedge (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (\\
& \forall X4.(m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow (\neg (X3 \in X2) \wedge ((X4 \in \\
& X2) \wedge ((k3_funct_2 (u1_struct_0 X0) (u1_struct_0 k3_topmetr) X1 \\
& X3 = k4_seq_4 (k1_weierstr (k7_reset_1 (u1_struct_0 X0) (u1_struct_0 \\
& k3_topmetr) X1 X2))) \wedge (k3_funct_2 (u1_struct_0 X0) (u1_struct_0 \\
& k3_topmetr) X1 X4 = k5_seq_4 (k1_weierstr (k7_reset_1 (u1_struct_0 \\
& X0) (u1_struct_0 k3_topmetr) X1 X2))))))))))))) \Rightarrow
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (\\
& k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (m1_subset_1 (k7_reset_1 \\
& X0 X1 X2 X3) (k1_zfmisc_1 X1))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0. (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 k3_topmetr))) \Rightarrow \\
& (k3_weierstr X0 = k5_seq_4 (k1_weierstr X0))
\end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (u1_struct_0 \\ & X0) (u1_struct_0 k3_topmetr)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 k3_topmetr)))))) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow \\ & (\neg (X2 \neq k1_xboole_0) \wedge ((v2_compts_1 X2 X0) \wedge ((v5_pre_topc X1 X0 \\ & k3_topmetr) \wedge (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (\\ & \neg (X3 \in X2) \wedge (k3_funct_2 (u1_struct_0 X0) (u1_struct_0 k3_topmetr) \\ & X1 X3 = k3_weierstr (k7_relset_1 (u1_struct_0 X0) (u1_struct_0 \\ & k3_topmetr) X1 X2)))))))))) \end{aligned}$$