

t15\_zf\_model  
(TMJfejwYoLkszFBtzfF62CNFQ3k4Q9PpQaT)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zf\_lang : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_zf\_lang : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_zf\_model : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_zf\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_zf\_model : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_zf\_lang X1) \wedge ( \\ & m2\_finseq\_1 X1 k5\_numbers)) \Rightarrow (\forall X2.((v1\_zf\_lang X2) \wedge (m2\_finseq\_1 \\ & X2 k5\_numbers)) \Rightarrow (\forall X3.((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 \\ & k1\_zf\_lang X0) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_zf\_lang \\ & X0)))))) \Rightarrow (((X3 \in k5\_zf\_model X1 X0) \wedge (X3 \in k5\_zf\_model X2 X0)) \Leftrightarrow (X3 \in \\ & k5\_zf\_model (k7\_zf\_lang X1 X2) X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1\_zf\_lang X0) \wedge (m1\_finseq\_1 X0 k5\_numbers)) \wedge \\ & ((v1\_zf\_lang X1) \wedge (m1\_finseq\_1 X1 k5\_numbers))) \Rightarrow (v1\_zf\_lang \\ & (k7\_zf\_lang X0 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_finseq\_1 X0 k5\_numbers) \wedge (m1\_finseq\_1 X1 k5\_numbers)) \Rightarrow (m2\_finseq\_1 (k7\_zf\_lang X0 X1) k5\_numbers) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 k1\_zf\_lang X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & k1\_zf\_lang X0)))))) \Rightarrow (\forall X2.((v1\_zf\_lang X2) \wedge (m2\_finseq\_1 \\ & X2 k5\_numbers)) \Rightarrow ((r1\_zf\_model X0 X1 X2) \Leftrightarrow (X1 \in k5\_zf\_model X2 X0)))) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 k1\_zf\_lang X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & k1\_zf\_lang X0)))))) \Rightarrow (\forall X2.((v1\_zf\_lang X2) \wedge (m2\_finseq\_1 \\ & X2 k5\_numbers)) \Rightarrow (\forall X3.((v1\_zf\_lang X3) \wedge (m2\_finseq\_1 X3 \\ & k5\_numbers)) \Rightarrow ((r1\_zf\_model X0 X1 (k7\_zf\_lang X2 X3)) \Leftrightarrow ((r1\_zf\_model \\ & X0 X1 X2) \wedge (r1\_zf\_model X0 X1 X3)))))) \end{aligned}$$