

t165_xreal_1

(TMHm8VvECw2EFBxDvNKvLRqdu1FypRYp26W)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 (k4_xcmplx_0 np_1) = k4_xcmplx_0 X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k4_xcmplx_0 (k4_xcmplx_0 np_1) = np_1 \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X2 k6_numbers)) \Rightarrow \\ & (r1_xxreal_0 (k3_xcmplx_0 X1 X2) (k3_xcmplx_0 X0 X2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 X1) (k4_xcmplx_0 X0)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k3_xcmplx_0 X0 X1)) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (13)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge ((r1_xxreal_0 (k4_xcmplx_0 np_1) X0) \wedge (r1_xxreal_0 (k4_xcmplx_0 np_1) X1))) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X1) np_1)))$$