

t165\_xxreal\_1

(TMK7ia7HeCLCxgZXEuRYwWcbBCJNEpRDCnR)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_membered : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_numbers : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ & (r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((X0 \in k1\_xxreal\_1 X1 X2) \Leftrightarrow ((r1\_xxreal\_0 X1 X0) \wedge \\ & (r1\_xxreal\_0 X0 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (r1\_xxreal\_0 X0 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v2\_membered X0) \wedge (v2\_membered X1)) \Rightarrow (v2\_membered (k2\_xboole\_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (v2\_membered (k1\_xxreal\_1 X0 X1)) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 = k2\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (k1\_xxreal\_1 \\ X0 X1 = ReplSep (toset (\lambda X2 : \iota.m1\_subset\_1 X2 k7\_numbers)) \\ (\lambda X2 : \iota.(r1\_xxreal\_0 X0 X2) \wedge (r1\_xxreal\_0 X2 X1)) (\lambda X2 : \\ \iota.X2)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2\_membered X0) \Rightarrow (\forall X1.(v2\_membered X1) \Rightarrow (( \\ X0 = X1) \Leftrightarrow (\forall X2.(v1\_xxreal\_0 X2) \Rightarrow ((X2 \in X0) \Leftrightarrow (X2 \in X1)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow ( \\ (r1\_xxreal\_0 X0 X1) \vee (r1\_xxreal\_0 X1 X0)) \end{aligned} \quad (9)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ (v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ (k2\_xboole\_0 (k1\_xxreal\_1 X0 X1) (k1\_xxreal\_1 X1 X2) = k1\_xxreal\_1 \\ X0 X2)))) \end{aligned}$$