

t16_bvfunc11

(TMJ5GjK5jJgq3HvHo4nwATM4ix2MdN2GmBC)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_2 : \iota \Rightarrow \iota$ be given. Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. (m1_subset_1 X2 \\
 & (k1_zfmisc_1 (k1_bvfunc_2 X0))) \Rightarrow (\forall X3. (m1_eqrel_1 X3 X0) \Rightarrow \\
 & (\forall X4. (m1_eqrel_1 X4 X0) \Rightarrow (r1_bvfunc_1 X0 (k6_bvfunc_2 X0 \\
 & X1 X2 X3) (k7_bvfunc_2 X0 X1 X2 X4))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (((r1_bvfunc_1 X0 X1 X2) \wedge (\\
 & r1_bvfunc_1 X0 X2 X1)) \Rightarrow (r2_funct_2 X0 k6_margrel1 X1 X2)) \wedge (((r1_bvfunc_1 \\
 & X0 X1 X2) \wedge (r1_bvfunc_1 X0 X2 X3)) \Rightarrow (r1_bvfunc_1 X0 X1 X3))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
& (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
& k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.(m1_subset_1 X2 \\
& (k1_zfmisc_1 (k1_bvfunc_2 X0))) \Rightarrow (\forall X3.(m1_eqrel_1 X3 X0) \Rightarrow \\
& (\forall X4.(m1_eqrel_1 X4 X0) \Rightarrow ((v2_bvfunc_2 X2 X0) \Rightarrow (r1_bvfunc_1 \\
& X0 (k7_bvfunc_2 X0 (k1_bvfunc_1 X0 (k6_bvfunc_2 X0 X1 X2 X3)) X2 X4) \\
& (k7_bvfunc_2 X0 (k7_bvfunc_2 X0 (k1_bvfunc_1 X0 X1) X2 X4) X2 X3)))))) \\
& \hspace{15em} (3)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\
& (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 \\
& X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((m1_subset_1 \\
& X2 (k1_zfmisc_1 (k1_bvfunc_2 X0))) \wedge (m1_eqrel_1 X3 X0))) \Rightarrow ((v1_funct_1 \\
& (k7_bvfunc_2 X0 X1 X2 X3)) \wedge ((v1_funct_2 (k7_bvfunc_2 X0 X1 X2 X3) \\
& X0 k6_margrel1) \wedge (m1_subset_1 (k7_bvfunc_2 X0 X1 X2 X3) (k1_zfmisc_1 \\
& (k2_zfmisc_1 X0 k6_margrel1)))))) \\
& \hspace{15em} (4)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\
& (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 \\
& X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((m1_subset_1 \\
& X2 (k1_zfmisc_1 (k1_bvfunc_2 X0))) \wedge (m1_eqrel_1 X3 X0))) \Rightarrow ((v1_funct_1 \\
& (k6_bvfunc_2 X0 X1 X2 X3)) \wedge ((v1_funct_2 (k6_bvfunc_2 X0 X1 X2 X3) \\
& X0 k6_margrel1) \wedge (m1_subset_1 (k6_bvfunc_2 X0 X1 X2 X3) (k1_zfmisc_1 \\
& (k2_zfmisc_1 X0 k6_margrel1)))))) \\
& \hspace{15em} (5)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\
& (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
& k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k1_bvfunc_1 \\
& X0 X1)) \wedge ((v1_funct_2 (k1_bvfunc_1 X0 X1) X0 k6_margrel1) \wedge (m1_subset_1 \\
& (k1_bvfunc_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \\
& \hspace{15em} (6)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
& (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
& k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.(m1_subset_1 X2 \\
& (k1_zfmisc_1 (k1_bvfunc_2 X0))) \Rightarrow (\forall X3.(m1_eqrel_1 X3 X0) \Rightarrow \\
& (\forall X4.(m1_eqrel_1 X4 X0) \Rightarrow ((v2_bvfunc_2 X2 X0) \Rightarrow (r1_bvfunc_1 \\
& X0 (k6_bvfunc_2 X0 (k1_bvfunc_1 X0 (k6_bvfunc_2 X0 X1 X2 X3)) X2 X4) \\
& (k7_bvfunc_2 X0 (k7_bvfunc_2 X0 (k1_bvfunc_1 X0 X1) X2 X4) X2 X3)))))) \\
& \hspace{15em} (6)
\end{aligned}$$