

# t16\_euclid (TMZPmVNwpoVBJGYMgYE- hBL5srCb6xgP13CG)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Let  $k12\_euclid : \iota \Rightarrow \iota$  be given. Let  $k8\_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k5\_euclid : \iota \Rightarrow \iota$  be given. Let  $k4\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_rvsum\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $m1\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k45\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_euclid : \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v2\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v4\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v5\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(m2\_finseq\_2 X1 k1\_numbers (k1\_euclid X0)) \Rightarrow ((k12\_euclid X1 = k6\_numbers) \Rightarrow (X1 = k5\_euclid X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (k12\_euclid (k5\_euclid X0) = k6\_numbers) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(m2\_finseq\_2 X1 k1\_numbers (k4\_finseq\_2 X0 k1\_numbers)) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 k1\_numbers (k4\_finseq\_2 X0 k1\_numbers)) \Rightarrow ((k9\_rvsum\_1 X0 X1 X2 = k5\_finseq\_2 k5\_numbers X0 k6\_numbers) \Rightarrow (X1 = X2)))) \quad (3)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0) \Rightarrow (\forall X1.(m2\_finseq\_2\ X1\ k1\_numbers\ (k4\_finseq\_2\ X0\ k1\_numbers)) \Rightarrow (k9\_rsum\_1\ X0\ X1\ X1 = k5\_finseq\_2\ k5\_numbers\ X0\ k6\_numbers)) \quad (4)$$

Assume the following.

$$m1\_subset\_1\ k1\_xboole\_0\ k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_2\ X1\ X0) \Rightarrow (\forall X2.(m2\_finseq\_2\ X2\ X0\ X1) \Leftrightarrow (m1\_subset\_1\ X2\ X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1\ X1\ X0) \Leftrightarrow (m1\_finseq\_1\ X1\ X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7\_ordinal1\ X0) \wedge ((m1\_subset\_1\ X1\ (k4\_finseq\_2\ X0\ k1\_numbers)) \wedge (m1\_subset\_1\ X2\ (k4\_finseq\_2\ X0\ k1\_numbers)))) \Rightarrow (k9\_rsum\_1\ X0\ X1\ X2 = k45\_valued\_1\ X1\ X2) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7\_ordinal1\ X0) \wedge ((m1\_subset\_1\ X1\ (k1\_euclid\ X0)) \wedge (m1\_subset\_1\ X2\ (k1\_euclid\ X0)))) \Rightarrow (k8\_euclid\ X0\ X1\ X2 = k45\_valued\_1\ X1\ X2) \quad (9)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (10)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0\ X0) \wedge ((v7\_ordinal1\ X1) \wedge (m1\_subset\_1\ X2\ X0))) \Rightarrow (k5\_finseq\_2\ X0\ X1\ X2 = k2\_finseq\_2\ X1\ X2) \quad (12)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0) \Rightarrow (k5\_euclid\ X0 = k4\_euclid\ X0) \quad (13)$$

Assume the following.

$$\begin{aligned} & \exists X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & k5\_numbers))) \wedge ((\neg v1\_xboole\_0 X0) \wedge ((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 \\ & X0 k5\_numbers) \wedge ((v5\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge \\ & ((v1\_partfun1 X0 k5\_numbers) \wedge ((v1\_funct\_2 X0 k5\_numbers k5\_numbers) \wedge \\ & ((v1\_valued\_0 X0) \wedge ((v2\_valued\_0 X0) \wedge ((v3\_valued\_0 X0) \wedge ((v4\_valued\_0 \\ & X0) \wedge (v5\_valued\_0 X0)))))))))) \end{aligned} \quad (14)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_2 X1 X0) \Rightarrow (\forall X2.(m2\_finseq\_2 X2 X0 X1) \Rightarrow (m2\_finseq\_1 X2 X0)) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)) \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v7\_ordinal1 X0) \wedge ((m1\_subset\_1 \\ & X1 (k1\_euclid X0)) \wedge (m1\_subset\_1 X2 (k1\_euclid X0)))) \Rightarrow (m2\_finseq\_2 \\ & (k8\_euclid X0 X1 X2) k1\_numbers (k1\_euclid X0)) \end{aligned} \quad (18)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (m2\_finseq\_2 (k5\_euclid X0) k1\_numbers (k1\_euclid X0)) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.(v7\_ordinal1 X0) \Rightarrow (m1\_finseq\_2 (k4\_finseq\_2 X0 X1) X1) \quad (20)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finseq\_1 X0) \wedge (v3\_valued\_0 X0)))) \Rightarrow (m1\_subset\_1 (k12\_euclid X0) k1\_numbers) \quad (21)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (k4\_euclid X0 = k5\_finseq\_2 k1\_numbers X0 k6\_numbers) \quad (22)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (k1\_euclid X0 = k4\_finseq\_2 X0 k1\_numbers) \quad (23)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_xboole\_0 X0)\Rightarrow(\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0))\Rightarrow(v1\_xboole\_0 X2)) \quad (24)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow(v5\_relat\_1 X1 X0) \quad (25)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge(v5\_relat\_1 X0 k1\_numbers))\Rightarrow((v1\_relat\_1 X0)\wedge(v3\_valued\_0 X0)) \quad (26)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(\forall X1.(m2\_finseq\_2 X1 k1\_numbers (k1\_euclid X0))\Rightarrow(\forall X2.(m2\_finseq\_2 X2 k1\_numbers (k1\_euclid X0))\Rightarrow((k12\_euclid (k8\_euclid X0 X1 X2) = k6\_numbers)\Leftrightarrow(X1 = X2))))$$