

t16_finseq_4
(TMHtR7tbUwfhSpF7zXmjnVxSRRB85rv7AjB)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k4_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (X1 = k1_funct_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (3)$$

Assume the following.

$$\forall X0. (v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (6)$$

Assume the following.

$$\forall X0.k5_finseq_1 X0 = k1_tarski (k4_tarski np_1 X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(X2 = X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (9)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(k7_partfun1 X0 (k12_finseq_1 X0 X1) np_1 = X1))$$