

t16_gr_cy_3 (TMd-
JqxGR5tDA4YNcUPDtwyBa5DCpuDKLK2J)

October 27, 2020

Let $k1_gr_cy_3 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_newton X0 k6_numbers = np_1) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (4)$$

Assume the following.

$$k6_xcmplx_0 np_1 np_1 = np_0 \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_newton X0 X1 = k1_newton X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(k1_gr_cy_3\ X0 = k6_xcmplx_0\ (k2_newton\ np_2\ X0)\ np_1) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0)\Rightarrow(v7_ordinal1\ X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0\ X0)\Rightarrow(v1_xcmplx_0\ X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \quad (11)$$

Theorem 1 $k1_gr_cy_3\ k6_numbers = k6_numbers$.