

t16_grfunc_1

(TMT2udsWRXVMGhvvdKVsyErwTaeq5NTEZY1)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k4_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (X1 = k1_funct_1 X2 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarski X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (\forall X3. ((v1_relat_1 X3) \wedge (v1_funct_1 X3)) \Rightarrow (\neg (X0 \in k9_xtuple_0 X1) \wedge ((X1 = k2_xboole_0 X2 X3) \wedge ((k1_funct_1 X1 X0 \neq k1_funct_1 X2 X0) \wedge (k1_funct_1 X1 X0 \neq k1_funct_1 X3 X0)))))))$$