

t16_irrat_1

(TMTrAb4j6hrV9g9ADh8EQzA87Lb77jeVgUX)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 k5_numbers) \Rightarrow (\forall X3.(m2_finseq_1 \\ & X3 X0) \Rightarrow (((r1_xxreal_0 np_1 X2) \wedge (r1_xxreal_0 (k2_nat_1 X2 X1) \\ & (k3_finseq_1 X3))) \Rightarrow (k1_funct_1 (k2_rfinseq X0 X1 X3) X2 = k1_funct_1 \\ & X3 (k2_nat_1 X2 X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers)\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow((\neg r1_xxreal_0 (k3_finseq_1 X1) X0)\Rightarrow((k2_nat_1 X0 np_1 \in k4_finseq_1 X1)\wedge(k1_funct_1 X1 (k2_nat_1 X0 np_1) \in k10_xtuple_0 X1)))) \quad (10)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (14)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k4_ordinal1) \Rightarrow (v7_ordinal1 \ X0) \quad (16)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ ((v1_relat_1 \ X0) \wedge ((v4_relat_1 \ X0 \ k5_numbers) \wedge ((v1_funct_1 \ X0) \wedge \\ (v1_finseq_1 \ X0)))) \quad (17) \end{aligned}$$

Assume the following.

$$\forall X0.(v1_xboole_0 \ X0) \Rightarrow (\forall X1.(m1_subset_1 \ X1 \ (k1_zfmisc_1 \ X0)) \Rightarrow (v1_xboole_0 \ X1)) \quad (18)$$

Theorem 1

$$\begin{aligned} \forall X0.(m2_subset_1 \ X0 \ k1_numbers \ k5_numbers) \Rightarrow (\forall X1. \\ (\neg v1_xboole_0 \ X1) \Rightarrow (\forall X2.(m2_finseq_1 \ X2 \ X1) \Rightarrow ((r1_xxreal_0 \\ np_1 \ X0) \Rightarrow ((r1_xxreal_0 \ (k3_finseq_1 \ X2) \ X0) \vee (k1_funct_1 \ (k2_rfinseq \\ X1 \ np_1 \ X2) \ X0 = k1_funct_1 \ X2 \ (k2_nat_1 \ X0 \ np_1)))))) \end{aligned}$$