

t16_latsum_1 (TM-
StmHKEYm7mNBKCaGHvxmz4iYJJMkpoC5P)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_orders_2 : \iota \Rightarrow o$ be given. Let $v4_orders_2 : \iota \Rightarrow o$ be given. Let $v5_orders_2 : \iota \Rightarrow o$ be given. Let $v1_lattice3 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $v1_waybel_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v12_waybel_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k13_lattice3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v5_orders_2 X0) \wedge ((v1_lattice3 X0) \wedge (l1_orders_2 \\ & X0))) \Rightarrow (\forall X1.((v12_waybel_0 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (u1_struct_0 X0)))) \Rightarrow ((v1_waybel_0 X1 X0) \Leftrightarrow (\forall X2.(m1_subset_1 \\ & X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 \\ & X0)) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k13_lattice3 X0 X2 X3 \in X1)))))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_orders_2 X0) \wedge ((v4_orders_2 \\ & X0) \wedge ((v5_orders_2 X0) \wedge ((v1_lattice3 X0) \wedge (l1_orders_2 X0)))))) \Rightarrow \\ & (\forall X1.((v1_waybel_0 X1 X0) \wedge ((v12_waybel_0 X1 X0) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (u1_struct_0 X0)))))) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 \\ & X0)) \Rightarrow (((X2 \in X1) \wedge (X3 \in X1)) \Rightarrow (k13_lattice3 X0 X2 X3 \in X1)))) \end{aligned}$$