

t16_matrix_2

(TMSMtG4u96qcyxEpgox9hbQVWBKW4BwVVF4)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_matrix_1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_2 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_matrix_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_matrix_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_matrix_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k13_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2. \\ & (\neg v1_xboole_0 X2) \Rightarrow (\forall X3.(m1_matrix_1 X3 X2 X0 X1) \Rightarrow (\forall X4. \\ & (v7_ordinal1 X4) \Rightarrow ((X4 \in k2_finseq_1 X0) \Rightarrow (k1_funct_1 X3 X4 = k8_matrix_1 \\ & X2 X3 X4)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\\ & \forall X2.((v1_matrix_1 X2) \wedge (m2_finseq_1 X2 (k3_finseq_2 X1))) \Rightarrow \\ & ((k3_finseq_1 X2 = X0) \Rightarrow (m1_matrix_1 X2 X1 X0 (k1_matrix_1 X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.k3_finseq_2 X0 = k13_finseq_1 X0 \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1) \wedge (v1_finseq_1 X1)) \quad (11)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v1_matrix_1 X0)))) \Rightarrow (m1_subset_1 (k1_matrix_1 X0) k5_numbers) \quad (14)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow ((X1 = k3_finseq_1 X0) \Leftrightarrow (k2_finseq_1 X1 = k9_xtuple_0 X0))) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (16)$$

Theorem 1

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_matrix_1 X1) \wedge \\ (m2_finseq_1 X1 (k3_finseq_2 X0))) \Rightarrow (\forall X2. (v7_ordinal1 \\ X2) \Rightarrow ((X2 \in k4_finseq_1 X1) \Rightarrow (k1_funct_1 X1 X2 = k8_matrix_1 X0 X1 \\ X2)))) \end{aligned}$$