

t16_ordinal1
(TMcE9zzkmyuhR27bSanfGb7af3RyJJUXnFa)

October 27, 2020

Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_ordinal1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow (\forall X1.(v3_ordinal1 X1) \Rightarrow (\neg (\neg X0 \in X1) \wedge ((X0 \neq X1) \wedge (\neg X1 \in X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v3_ordinal1 X0) \wedge (v3_ordinal1 X1)) \Rightarrow ((r1_ordinal1 X0 X1) \Leftrightarrow (r1_tarski X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_ordinal1 X0) \Leftrightarrow (\forall X1.(X1 \in X0) \Rightarrow (r1_tarski X1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1_tarski X0 X1) \wedge (r1_tarski X1 X0)) \quad (4)$$

Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow ((v1_ordinal1 X0) \wedge (v2_ordinal1 X0)) \quad (5)$$

Theorem 1

$$\forall X0.(v3_ordinal1 X0) \Rightarrow (\forall X1.(v3_ordinal1 X1) \Rightarrow ((r1_ordinal1 X0 X1) \vee (X1 \in X0)))$$