

t16\_polyform  
(TMaV71zhgYnmMpgr5AEfFVydWV1r7wVnobf)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (k3\_finseq\_1 (k7\_finseq\_1 X0 X1) = k2\_nat\_1 (k3\_finseq\_1 X0) (k3\_finseq\_1 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (4)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \wedge ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow \\ & ((v1\_relat\_1 (k7\_finseq\_1 X0 X1)) \wedge ((v1\_funct\_1 (k7\_finseq\_1 X0 X1)) \wedge (v1\_finseq\_1 (k7\_finseq\_1 X0 X1)))) \end{aligned} \quad (6)$$

Assume the following.

$$m1\_subset\_1\ k5\_numbers\ (k1\_zfmisc\_1\ k1\_numbers) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1\ X0)\wedge((v1\_funct\_1\ X0)\wedge(v1\_finseq\_1\ X0)))\Rightarrow \\ (m2\_subset\_1\ (k3\_finseq\_1\ X0)\ k1\_numbers\ k5\_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1\ X0\ k4\_ordinal1)\Rightarrow(v7\_ordinal1\ X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ X0))\Rightarrow(v1\_xboole\_0\ X1)) \quad (10)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1\ X0)\wedge((v1\_funct\_1\ X0)\wedge(v1\_finseq\_1\ X0)))\Rightarrow \\ (\forall X1.((v1\_relat\_1\ X1)\wedge((v1\_funct\_1\ X1)\wedge(v1\_finseq\_1\ X1)))\Rightarrow(\forall X2.((v1\_relat\_1\ X2)\wedge((v1\_funct\_1\ X2)\wedge(v1\_finseq\_1\ X2)))\Rightarrow(k3\_finseq\_1\ (k7\_finseq\_1\ (k7\_finseq\_1\ X0\ X1)\ X2) = k2\_xcmplx\_0 \\ (k2\_xcmplx\_0\ (k3\_finseq\_1\ X0)\ (k3\_finseq\_1\ X1))\ (k3\_finseq\_1\ X2))))$$