

t16\_subset\_1  
 (TMJs1tFh25ogpo6QCvnnzk6nL4sDDxNpTCd)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (\forall X2. \\ & (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow ((r1\_tarski X1 X2) \Leftrightarrow (r1\_tarski \\ & (k3\_subset\_1 X0 X2) (k3\_subset\_1 X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (k3\_subset\_1 X0 (k3\_subset\_1 X0 X1) = X1) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (m1\_subset\_1 (k3\_subset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \tag{3}$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (\forall X2. \\ & (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow ((r1\_tarski X1 (k3\_subset\_1 \\ & X0 X2)) \Rightarrow (r1\_tarski X2 (k3\_subset\_1 X0 X1)))) \end{aligned}$$