

t16_tops_2 (TMbRUFcUgKPvxRVhmkEN- qyT3LNVdg5yYWVz)

October 27, 2020

Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_tops_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k4_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \tag{3}$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow ((v2_tops_2 X1 X0) \Leftrightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((X2 \in X1) \Rightarrow (v4_pre_topc X2 X0)))))) \tag{4}$$

Theorem 1

$$\forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (((v2_tops_2 X1 X0) \wedge (v2_tops_2 X2 X0)) \Rightarrow (v2_tops_2 (k4_subset_1 (k1_zfmisc_1 (u1_struct_0 X0)) X1 X2) X0))))))$$