

t16\_xxreal\_0 (TMN-  
VEiAF3nrv655o8JMUo6HwkTXzNcWh28)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k4\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ (r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X1 \\ X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X1)))) \end{aligned} \tag{1}$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ k4\_xxreal\_0 X0 X1 = X0) \vee (k4\_xxreal\_0 X0 X1 = X1)))$$