

t172_glib_001 (TMXN- jqvG8tRYRU2NsaDrVz5BUCQEuhnt2XU)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $m3_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_000 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k14_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k24_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k25_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_glib_000 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. \forall X2. \\ (m2_glib_000 X2 X0 (k6_glib_000 X0) (k6_subset_1 (k7_glib_000 \\ X0) X1)) \Rightarrow ((k24_glib_000 X0 X2 = k6_glib_000 X0) \wedge (k25_glib_000 \\ X0 X2 = k6_subset_1 (k7_glib_000 X0) X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. (m1_glib_000 \\ X1 X0) \Rightarrow (\forall X2. (m3_glib_001 X2 X0) \Rightarrow (((r1_tarski (k13_glib_001 \\ X0 X2) (k24_glib_000 X0 X1)) \wedge (r1_tarski (k14_glib_001 X0 X2) (k25_glib_000 \\ X0 X1))) \Rightarrow (m3_glib_001 X2 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ & ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \wedge (m1_glib_000 \\ & X1 X0)) \Rightarrow (k25_glib_000 X0 X1 = k7_glib_000 X1) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_relat_1 X0) \wedge ((v4_relat_1 \\ & X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 \\ & X0)))))) \Rightarrow (\forall X3. (m2_glib_000 X3 X0 X1 X2) \Rightarrow (m1_glib_000 X3 \\ & X0)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ & ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \wedge (m3_glib_001 \\ & X1 X0)) \Rightarrow ((v1_finset_1 (k14_glib_001 X0 X1)) \wedge (m1_subset_1 (k14_glib_001 \\ & X0 X1) (k1_zfmisc_1 (k7_glib_000 X0)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ & ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \wedge (m3_glib_001 \\ & X1 X0)) \Rightarrow ((v1_finset_1 (k13_glib_001 X0 X1)) \wedge (m1_subset_1 (k13_glib_001 \\ & X0 X1) (k1_zfmisc_1 (k6_glib_000 X0)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow \\ & (X2 \in X1)) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow \\ & (X2 = X0)) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. (m3_glib_001 \\ & X1 X0) \Rightarrow (\forall X2. \forall X3. (m2_glib_000 X3 X0 (k6_glib_000 \\ & X0) (k6_subset_1 (k7_glib_000 X0) (k1_tarski X2))) \Rightarrow ((\neg X2 \in k14_glib_001 \\ & X0 X1) \Rightarrow (m3_glib_001 X1 X3)))) \end{aligned}$$