

t176_relat_1
(TML1xrx6VrZZCLsvvYazH1nNdjZvuxKQ7Th)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow (k5_relat_1 X2 (k6_subset_1 X0 X1) = k6_subset_1 (k5_relat_1 X2 X0) (k5_relat_1 X2 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k9_xtuple_0 X1) X0) \Rightarrow (k5_relat_1 X1 X0 = X1)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((r1_tarski (k9_xtuple_0 X2) X0) \Rightarrow (k6_subset_1 X2 (k5_relat_1 X2 X1) = k5_relat_1 X2 (k6_subset_1 X0 X1)))$$