

t17_bvfunc26

(TMZ1VeyiW13VCGA4d9P1CFkjhAh8LihTbgj)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_bvfunc26 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge (\\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (\\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge (\\ & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (\\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc26 X0 X1 (k5_bvfunc_1 X0 X2 X3)) (k2_bvfunc_1 X0 (k1_bvfunc_1 \\ & X0 (k2_bvfunc_1 X0 X1 X2)) (k1_bvfunc_1 X0 (k2_bvfunc_1 X0 X1 X3)))))) \\ & \tag{1} \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow (k2_bvfunc_1 X0 X1 X1 = X1) \\ & \tag{2} \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge (\\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (\\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc26 X0 X1 (k5_bvfunc_1 X0 X1 X2)) (k2_bvfunc_1 X0 (k1_bvfunc_1 \\ & X0 X1) (k1_bvfunc_1 X0 (k2_bvfunc_1 X0 X1 X2)))))) \end{aligned}$$